

Allergies: See Allergy Sheet	ACTION
Peripherally Inserted Central Catheter (PICC) Pre and Post Proced Order Set	ure
Information Sheet	
\square The PICC patient information sheet has been provided to the patient by the referring provider signed and brought to the PICC team the day of the procedure	to be
Laboratory Investigations	
For patients on therapeutic anticoagulation:	
☑ INR within 7 days of procedure if patient currently on warfarin☐ Resume (previously order therapeutic anticoagulation) hours post	insertion
For patients on prophylactic anticoagulation:	
 ☑ Do not hold anticoagulation for VTE prophylaxis ☑ Do not hold acetylsalicylic acid or clopidogrel or other anti-platelet therapy 	
Pain Management	
☐ Have available Lidocaine 1% for intradermal injection at insertion site	
Post procedure Diagnostic Imaging ☐ Sherlock 3CG Bard Technology (attach printed tracing to space provided at end of set) ☐ CXR-indication: PICC tip placement, unable to confirm using Sherlock 3CG Bard technology ☐ CXR-indication: PICC tip placement, alteration of cardiac rhythm post placement	
Post PICC Insertion monitoring	
 ☑ Inspect site q15 minutes x2, and prior to discharge and report suspected complications or con ☑ Notify MRP immediately for local bleeding, respiratory distress, suspected catheter migration, catheter-borne infection 	
PICC Line Maintenance	
 ☐ Flush PICC with 20mL 0.9% sodium chloride with a needleless valve ☐ Do not start infusion until tip placement is confirmed ☐ Use infusion pump for all infusions 	
Practitioner's Signature Printed Name	
Date/Time	



Allergies: See Allergy Sheet	ACTION
Peripherally Inserted Central Catheter (PICC) Pre and Post Procedure Order Set	
Dressing	
 ☑ Apply gauze at exit site and cover with transparent dressing ☑ Remove transparent dressing and gauze from exit site at 24 hours post insertion ☑ Reapply gauze only if oozing noted at exit site, cover with transparent dressing ☑ Change PICC dressing and securement device q 7 days and PRN, if gauze used at the exit site, change dressing q 48 hours to ensure inspection of the site 	
Flushing	
 ☑ Flush each lumen with 10mL 0.9% sodium chloride after each use, weekly if not used ☑ Flush lumen with 20mL 0.9% sodium chloride after any blood sampling, viscous solutions (blood products, CT dye, TPN) ☑ Do not use a syringe smaller than 10mL ☑ If out-patient or being discharged prior to 24hours, flush each lumen with 10mL 0.9% sodium chloride, and arrange for 24 hour PICC line assessment and dressing change 	
Discharge	
 ☑ Provide PICC Line information if patient discharged home with PICC ☑ Discharge Referral to LIHN Home and Community Care for PICC Line care 	
Attach printed confirmation from Shelock 3CG-Tip confirmation system below:	
Post PICC Line Insertion Documentation	
Insertion site: Left Right Location of catheter tip: SVC Other: Location confirmed by: Internal measurement (cut length): cm External measurement (exit site to bifurcation): cm	
Practitioner's Signature Printed Name Date/Time	