



PEDIATRIC – IV Infusion Chart – norepinephrine 0.4 mg

norepinephrine 0.4 mg in 50 mL (0.008 mg/mL = 8 mcg/mL) SYRINGE Patient weight 1 kg or less

PREPARATION: Draw up 0.4 mL of norepinephrine 1 mg/mL
Draw up 49.6 mL of dextrose 5% into a 50 mL syringe
and add 0.4 mL of norepinephrine to the same 50 mL
syringe

CONCENTRATION: 8 mcg/mL

VOLUME: 50 mL

PRESCRIBING UNITS: mcg/kg/min

INFUSION: 0.05 – 0.1 mcg/kg/min

DOSE (mcg/kg/min)	WEIGHT (kg)								
	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
RATE (mL/hour)									
0.05	0.08	0.11	0.15	0.19	0.23	0.26	0.3	0.34	0.38
0.06	0.09	0.14	0.18	0.23	0.27	0.32	0.36	0.41	0.45
0.07	0.11	0.16	0.21	0.26	0.32	0.37	0.42	0.47	0.53
0.08	0.12	0.18	0.24	0.3	0.36	0.42	0.48	0.54	0.6
0.09	0.14	0.2	0.27	0.34	0.41	0.47	0.54	0.61	0.68
0.1	0.15	0.23	0.3	0.38	0.45	0.53	0.6	0.68	0.75